

Memo

LET WEL:


- Indien 'n kandidaat 'n vraag TWEE KEER beantwoord, merk slegs die EERSTE poging.
- Volgehoue akkuraatheid is op ALLE aspekte van die nasienriglyne van toepassing.

QUESTION/VRAAG 1

1.1.1	$x^2 + 9x + 14 = 0$ $(x+7)(x+2) = 0$ $x = -7 \text{ or } x = -2$	✓ factors ✓ $x = -7$ ✓ $x = -2$ (3)
1.1.2	$4x^2 + 9x - 3 = 0$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $= \frac{-9 \pm \sqrt{9^2 - 4(4)(-3)}}{2(4)}$ $= \frac{-9 \pm \sqrt{129}}{8}$ $x = 0,29 \text{ or } x = -2,54$ <p>OR/OF</p> $x^2 + \frac{9}{4}x + \frac{81}{64} = \frac{3}{4} + \frac{81}{64}$ $\left(x + \frac{9}{8}\right)^2 = \frac{129}{64}$ $x + \frac{9}{8} = \pm \frac{\sqrt{129}}{8}$ $x = \frac{-9 \pm \sqrt{129}}{8}$ $x = 0,29 \text{ or } x = -2,54$	✓ substitution ✓ simplification ✓ $x = 0,29$ ✓ $x = -2,54$ <p>OR/OF</p> ✓ for adding $\frac{81}{64}$ on both sides ✓ simplification ✓ $x = 0,29$ ✓ $x = -2,54$ (4)
1.1.3	$\sqrt{x^2 - 5} = 2\sqrt{x}$ $x^2 - 5 = 4x$ $x^2 - 4x - 5 = 0$ $(x-5)(x+1) = 0$ $x = 5 \text{ or } x = -1$ $x = 5$	✓ $x^2 - 5 = 4x$ ✓ standard form ✓ both answers ✓ select $x = 5$ (4)

1.2	$ \begin{aligned} 3x - y &= 4 \\ y &= 3x - 4 \\ x^2 + 2xy - y^2 &= -2 \\ x^2 + 2x(3x - 4) - (3x - 4)^2 &= -2 \\ x^2 + 6x^2 - 8x - (9x^2 - 24x + 16) &= -2 \\ 7x^2 - 8x - 9x^2 + 24x - 16 &= -2 \\ -2x^2 + 16x - 14 &= 0 \\ x^2 - 8x + 7 &= 0 \\ (x - 7)(x - 1) &= 0 \\ x = 1 &\quad \text{or} \quad x = 7 \\ y = 3(1) - 4 &\quad y = 3(7) - 4 \\ y = -1 &\quad \text{or} \quad y = 17 \end{aligned} $ <p>OR/OF</p> $ \begin{aligned} 3x - y &= 4 \\ x &= \frac{y + 4}{3} \\ x^2 + 2xy - y^2 &= -2 \\ x^2 + 2xy - y^2 &= -2 \\ \left(\frac{y+4}{3}\right)^2 + 2\left(\frac{y+4}{3}\right)y - y^2 &= -2 \\ y^2 + 8y + 16 + 6y^2 + 24y - 9y^2 &= -18 \\ -2y^2 + 32y + 34 &= 0 \\ y^2 - 16y - 17 &= 0 \\ (y - 17)(y + 1) &= 0 \\ y = -1 &\quad \text{or} \quad y = 17 \\ x = \frac{-1+4}{3} &\quad x = \frac{17+4}{3} \\ x = 1 &\quad \text{or} \quad x = 7 \end{aligned} $	<ul style="list-style-type: none"> ✓ y subject of formula ✓ substitution ✓ correct standard form ✓ factors ✓ x-values ✓ y-values <p>OR/OF</p> <ul style="list-style-type: none"> ✓ x subject of formula ✓ substitution ✓ correct standard form ✓ factors ✓ y-values ✓ x-values
1.3.1	$ \begin{aligned} x^2 + 8x + 16 &> 0 \\ (x + 4)(x + 4) &> 0 \\ x \in R, x \neq -4 &\quad \text{or} \\ x \in (-\infty; -4) \text{ or } x \in (-4; \infty) &\quad \text{or} \\ x < -4 \text{ or } x > -4 & \end{aligned} $ <p>OR/OF</p> $ \begin{aligned} x^2 + 8x + 16 &> 0 \\ (x + 4)(x + 4) &> 0 \end{aligned} $ <p>The function values remain positive $x \in R, x \neq -4$</p>	<ul style="list-style-type: none"> ✓ $(x + 4)(x + 4)$ ✓✓ any one of the solutions <p>OR/OF</p> <ul style="list-style-type: none"> ✓ $(x + 4)(x + 4)$ ✓✓ any one of the solutions

1.3.2



For two negative unequal roots:
 $0 < p < 16$

OR/OF

$$x^2 + 8x + 16 = p$$

$$x^2 + 8x + 16 - p = 0$$

$$0 < 16 - p < 16$$

$$-16 < -p < 0$$

$$0 < p < 16$$

OR/OF

$$x^2 + 8x + 16 - p = 0$$

$$x = \frac{-8 \pm \sqrt{64 - 4(16 - p)}}{2}$$

$$0 < 64 - 4(16 - p) < 64$$

$$0 < 4p < 64$$

$$0 < p < 16$$

OR/OF

$$x^2 + 8x + 16 = p$$

$$x^2 + 8x + 16 - p = 0$$

Roots are real and unequal :

$$8^2 - 4(16 - p) > 0$$

$$4p > 0$$

$$p > 0$$

$$\text{Roots are : } \frac{-8 \pm \sqrt{4p}}{2}$$

For both roots to be negative :

$$\sqrt{4p} < 8$$

$$4p < 64$$

$$p < 16$$

$$0 < p < 16$$

- ✓ 0
- ✓ 16

- ✓ ✓ $0 < p < 16$ (4)

OR/OF


- ✓ 0
- ✓ 16

- ✓ ✓ $0 < p < 16$ (4)

- ✓ 0
- ✓ 16

- ✓ ✓ $0 < p < 16$ (4)

QUESTION/VRAAG 2

2.1.1  first differences: -9; -15; -21 second difference = -6	✓ first differences ✓ -6 (2)
2.1.2 $T_n = an^2 + bn + c$ $a = \frac{\text{second difference}}{2} = -3$ $3a + b = -9$ $3(-3) + b = -9$ $b = 0$ $a + b + c = 5$ $-3 + 0 + c = 5$ $c = 8$ $T_n = -3n^2 + 8$ OR/OF $T_n = T_1 + (n-1)d_1 + \frac{(n-1)(n-2)d_2}{2}$ $= 5 + (n-1)(-9) + \frac{(n-1)(n-2)(-6)}{2}$ $= 5 - 9n + 9 - 3n^2 + 9n - 6$ $T_n = -3n^2 + 8$	✓ $a = -3$ ✓ $b = 0$ ✓ $c = 8$ ✓ $T_n = -3n^2 + 8$ OR/OF ✓ $a = -3$ ✓ $b = 0$ ✓ $c = 8$ ✓ $T_n = -3n^2 + 8$ (4)
2.1.3 $-3n^2 + 8 = -25\ 939$ $-3n^2 = -25\ 947$ $n^2 = 8649$ $n = -93 \text{ or } n = 93$ <p>The 93rd term has a value of -25 939</p>	✓ $T_n = -25\ 939$ ✓ $n^2 = 8649$ ✓ answer (3)

2.2.1	$2k - 7 ; k + 8 \text{ and } 2k - 1$ $k + 8 - (2k - 7) = 2k - 1 - (k + 8)$ $-k + 15 = k - 9$ $2k = 24$ $k = 12$ $2k - 7; k + 8 \text{ and } 2k - 1$ $17; 20; 23 \dots$ $d = 3$ $T_{15} = 17 + 14(3)$ $= 59$	✓ $k + 8 - (2k - 7) = 2k - 1 - (k + 8)$ ✓ $k = 12$ ✓ 17 ✓ $d = 3$ ✓ $T_{15} = 59$ (5)
2.2.2	Sequence is 17 ; 20 ; 23 ; 26 ; 29 ; 32 Every alternate term of the sequence will be even / <i>Elke tweede term van die ry sal ewe wees</i> $20 + 26 + 32 + \dots$ $S_{30} = \frac{30}{2} [2(20) + (29)(6)]$ $= 15[40 + 174]$ $= 3210$ OR/OF $T_{30} = 20 + 29(6)$ $= 94$ $S_{30} = \frac{30}{2} (20 + 194)$ $= 3210$	✓ $20 + 26 + 32 + \dots$ ✓ $a = 20 \ d = 6$ ✓ subst into correct formula ✓ answer (4) ✓ $a = 20 \ d = 6$ ✓ $T_{30} = 94$ ✓ $S_{30} = \frac{30}{2} (20 + 194)$ ✓ answer (4) [18]

QUESTION/VRAAG 3

<p>3.1</p> $a + ar = 2$ $a(1+r) = 2$ $a = \frac{2}{1+r}$ <p>OR/OF</p> $\frac{a}{1-r} - 2 = \frac{1}{4}$ $4a - 8(1-r) = 1-r$ $4a - 8 + 8r = 1-r$ $4a = 9 - 9r$ $a = \frac{9-9r}{4}$ <p>OR/OF</p> $S_n = \frac{a(r^n - 1)}{r-1}$ $2 = \frac{a(r^2 - 1)}{r-1}$ $2 = \frac{a(r-1)(r+1)}{r-1}$ $2 = a(r+1)$ $a = \frac{2}{r+1}$ <p>OR/OF</p> $\frac{ar^2}{1-r} = \frac{1}{4}$ $a = \frac{1-r}{4r^2}$	<p>$\checkmark a + ar = 2$</p> $\checkmark a = \frac{2}{1+r}$ <p>OR/OF</p> $\checkmark \frac{a}{1-r} - 2 = \frac{1}{4}$ $\checkmark a = \frac{9-9r}{4}$ <p>OR/OF</p> $\checkmark 2 = \frac{a(r^2 - 1)}{r-1}$ $\checkmark a = \frac{2}{1+r}$ <p>OR/OF</p> $\checkmark \frac{ar^2}{1-r} = \frac{1}{4}$ $\checkmark a = \frac{1-r}{4r^2}$
--	--

<p>3.2</p> $S_{\infty} = T_1 + T_2 + \sum_{n=3}^{\infty} T_n$ $S_{\infty} = 2 + \frac{1}{4}$ $\frac{a}{1-r} = 2 + \frac{1}{4}$ $\frac{a}{1-r} = \frac{9}{4}$ $\left(\frac{2}{1+r}\right) \times \left(\frac{1}{1-r}\right) = \frac{9}{4}$ $\frac{2}{1-r^2} = \frac{9}{4}$ $8 = 9 - 9r^2$ $9r^2 = 1$ $r = \frac{1}{3}$ $a = \frac{3}{2}$	$\checkmark S_{\infty} = 2 + \frac{1}{4}$ $\checkmark \frac{a}{1-r} = \frac{9}{4}$ \checkmark substitution of a into the correct formula
<p>OR/OF</p> $S_{\infty} = T_1 + T_2 + \sum_{n=3}^{\infty} T_n$ $S_{\infty} = 2 + \frac{1}{4}$ $\frac{a}{1-r} = 2 + \frac{1}{4}$ $\frac{a}{1-r} = \frac{9}{4}$ $4a = 9 - 9r$ $r = \frac{9-4a}{9}$ $a + a\left(\frac{9-4a}{9}\right) = 2$ $9a + 9a - 4a^2 = 18$ $2a^2 - 9a + 9 = 0$ $(a-3)(2a-3) = 0$ $a = \frac{3}{2} \quad \text{or} \quad a = 3$ $r = \frac{1}{3} \quad \text{or} \quad r = -\frac{1}{3}$ <p style="text-align: center;">N/A</p>	<p>OR/OF</p> $\checkmark S_{\infty} = 2 + \frac{1}{4}$ $\checkmark \frac{a}{1-r} = \frac{9}{4}$ $\checkmark r = \frac{9-4a}{9}$ \checkmark substitution of a into the correct formula

	<p>OR/OF</p> $r = \frac{2-a}{a}$ $\frac{ar^2}{1-r} = \frac{1}{4}$ $4ar^2 = 1-r$ $4a\left(\frac{2-a}{a}\right)^2 = 1 - \frac{2-a}{a}$ $16 - 16a + 4a^2 = 2a + 2$ $2a^2 - 9a + 9 = 0$ $(2a-3)(a-3) = 0$ $a = \frac{3}{2} \quad a \neq 3$ $r = \frac{1}{3} \quad r \neq -\frac{1}{3}$ <p>OR/OF</p> $S_{\infty} = T_1 + T_2 + \sum_{n=3}^{\infty} T_n$ $S_{\infty} = 2 + \frac{1}{4}$ $\frac{a}{1-r} = 2 + \frac{1}{4}$ $\frac{a}{1-r} = \frac{9}{4}$ $\left(\frac{1-r}{4r^2}\right) \times \left(\frac{1}{1-r}\right) = \frac{9}{4}$ $\frac{1}{4r^2} = \frac{9}{4}$ $4 = 36r^2$ $9r^2 = 1$ $r = \frac{1}{3}$ $a = \frac{3}{2}$	<p>$\checkmark \quad r = \frac{1}{3}$</p> <p>OR/OF</p> <p>$\checkmark \quad r = \frac{2-a}{a}$</p> <p>$\checkmark \quad \frac{ar^2}{1-r} = \frac{1}{4}$</p> <p>$\checkmark \text{ substitution of } a$</p> <p>$\checkmark \quad (2a-3)(a-3) = 0$</p> <p>$\checkmark \quad a = \frac{3}{2}$</p> <p>$\checkmark \quad r = \frac{1}{3}$</p> <p>OR/OF</p> <p>$\checkmark \quad S_{\infty} = 2 + \frac{1}{4}$</p> <p>$\checkmark \quad \frac{a}{1-r} = \frac{9}{4}$</p> <p>$\checkmark \text{ substitution of } a$</p> <p>$\checkmark \quad 9r^2 = 1$</p> <p>$\checkmark \quad r = \frac{1}{3}$</p> <p>$\checkmark \quad a = \frac{3}{2}$</p>	(6)
			[8]

QUESTION/VRAAG 4

4.1 $f(x) = -ax^2 + bx + 6$ $f'(x) = -2ax + b$ $-2ax + b = 3$ $\text{at } x = -1$ $2a + b = 3 \quad [1]$ $f(-1) = \frac{7}{2}$ $-a - b + 6 = \frac{7}{2}$ $-2a - 2b + 12 = 7$ $2a + 2b = 5 \quad [2]$ $[2] - [1]$ $b = 2$ $2a + 2 = 3$ $a = \frac{1}{2}$ <p>OR/OF</p> $f'(x) = -2ax + b$ $3 = 2a + b$ $b = 3 - 2a$ $\frac{7}{2} = -a(-1)^2 + (3 - 2a)(-1) + 6$ $a + 3 = \frac{7}{2}$ $a = \frac{1}{2}$ $b = 2$	$\checkmark -2ax + b$ $\checkmark \checkmark 2a + b = 3$ $\checkmark -a - b + 6 = \frac{7}{2}$ $\checkmark \text{solve simultaneously}$ (5)
4.2 $f(x) = -\frac{1}{2}x^2 + 2x + 6$ <p>x-intercepts:</p> $-\frac{1}{2}x^2 + 2x + 6 = 0$ $-x^2 + 4x + 12 = 0$ $x^2 - 4x - 12 = 0$ $(x - 6)(x + 2) = 0$ $(-2; 0) \quad (6; 0)$	$\checkmark -\frac{1}{2}x^2 + 2x + 6 = 0$ $\checkmark (-2; 0)$ $\checkmark (6; 0)$ (3)

4.3	$f(x) = -\frac{1}{2}x^2 + 2x + 6$ $f'(x) = 0 \quad \text{or} \quad x = -\frac{b}{2a} \quad \text{or} \quad x = \frac{-2+6}{2}$ $-x + 2 = 0 \quad x = -\frac{2}{2\left(-\frac{1}{2}\right)} \quad x = 2$ $x = 2 \quad x = 2$ $y = -\frac{1}{2}(2)^2 + 2(2) + 6$ $= -2 + 4 + 6$ $= 8$ $\text{TP}(2; 8)$ OR/OF $y = -\frac{1}{2}(x^2 - 4x - 12)$ $= -\frac{1}{2}[(x-2)^2 - 4 - 12]$ $= -\frac{1}{2}(x-2)^2 + 8$ $\text{TP}(2; 8)$	$\checkmark -x + 2 / -\frac{2}{2\left(-\frac{1}{2}\right)} /$ $\frac{-2+6}{2}$ $\checkmark x = 2$ $\checkmark y = 8$ OR/OF $\checkmark -\frac{1}{2}(x-2)^2 + 8$ $\checkmark x = 2$ $\checkmark y = 8$ (3)
4.4 4.6	<p>The graph shows a parabola f opening downwards and a line g. The parabola f passes through the points $(-2, 0)$, $(0, 6)$, $(2, 8)$, and $(6, 0)$. The line g passes through the points $(-2, 0)$ and $(0, -1)$.</p>	4.4: f : \checkmark shape \checkmark x - intercepts \checkmark y - intercept $\checkmark (2; 8)$ (4) 4.6: g : \checkmark x - intercept \checkmark y - intercept (2)
4.5	$0 < x < 4$ or $(0; 4)$	$\checkmark 4$ $\checkmark \checkmark 0 < x < 4$ (3)
4.7	$x \leq -2$ or $-1 \leq x \leq 6$ OR/OF $(-\infty; -2] \text{ or } [-1; 6]$	$\checkmark x \leq -2$ $\checkmark \checkmark -1 \leq x \leq 6$ (3) [23]

QUESTION/VRAAG 5

5.1	$y \in R; y \neq -1$ OR/OF $y < -1$ or $y > -1$ OR/OF $y \in (-\infty; -1)$ or $y \in (-1; \infty)$ OR/OF $R - \{-1\}$	$\checkmark \checkmark$ answer (2)
5.2	$D(2; -1)$ $g(x) = \frac{2}{x-2} - 1$	$\checkmark D(2; -1)$ $\checkmark \frac{2}{x-2} - 1$ (2)
5.3	$f(x) = \log_3 x.$ $\log_3 t = 1$ OR/OF $g(x) = \frac{2}{x-2} - 1$ $t = 3$ $1 = \frac{2}{t-2} - 1$ $2 = \frac{2}{t-2}$ $2t - 4 = 2$ $t = 3$	\checkmark correct substitution of A $\checkmark \checkmark t = 3$ (3)
5.4	$x = \log_3 y$ $y = 3^x$	\checkmark interchange x and y $\checkmark y = 3^x$ (2)
5.5	$3^x < 3^1$ $x < 1$ OR/OF $3^x < 3^1$ $x \in (-\infty; 1)$	$\checkmark 3^x < 3^1$ $\checkmark x < 1$ (2) $\checkmark 3^x < 3^1$ $\checkmark x \in (-\infty; 1)$ (2)
5.6	Equation of the axis of symmetry: $y = -x + 1$ x -intercept of the axis of symmetry is at $x = 1$ f has an x -intercept at $B(1; 0)$ which is the same as the x -intercept of the axis of symmetry Point of intersection: $B(1; 0)$ OR/OF Since $BE = ED = 1$ and D lies on the axis of symmetry and the gradient of the axis of symmetry is -1 , B will also lie on the axis of symmetry. But B also lies on f . Therefore $B(1; 0)$ is the point of intersection between f and the axis of symmetry with a negative gradient./ <i>Omdat BE = ED = 1 en D op die simmetrie-as lê en die simmetrie-as se gradiënt -1 is, sal B ook op die simmetrie-as lê. Maar B lê ook op f. Dus is B(1; 0) die snypunt van f en die simmetrie-as met negatiewe gradiënt.</i>	$\checkmark \checkmark$ equation of axis of symmetry $\checkmark B$ or $(1; 0)$ OR/OF $\checkmark \checkmark BE = ED = 1$ $\checkmark B$ or $(1; 0)$ (3) [14]

QUESTION/VRAAG 6


6.1	$A = P(1+i)^n$ $12\ 146,72 = 10\ 000 \left(1 + \frac{r}{12}\right)^{36}$ $\left(1 + \frac{r}{12}\right)^{36} = 1,214672$ $1 + \frac{r}{12} = \sqrt[36]{1,214672}$ $= 1,005416$ $\frac{r}{12} = 0,005416$ $r = 0,06500$ $r = 6,5\%$	✓ $\frac{r}{12}$ ✓ $n = 36$ ✓ correct substitution into formula ✓ $1 + \frac{r}{12} = \sqrt[36]{1,214672}$ ✓ 6,5% (5)
6.2.1	$P = \frac{x \left[1 - (1+i)^{-n} \right]}{i}$ $235\ 000 = \frac{x \left[1 - \left(1 + \frac{0,11}{12} \right)^{-54} \right]}{\frac{0,11}{12}}$ $x = \frac{235\ 000 \times \frac{0,11}{12}}{\left[1 - \left(1 + \frac{0,11}{12} \right)^{-54} \right]}$ $= R5\ 536,95$ <p>His monthly instalment is R 5 536,95</p>	✓ $i = \frac{0,11}{12}$ ✓ $n = 54$ ✓ correct substitution in P ✓ answer (4)
6.2.2	Amount paid for the year : $(5\ 536,95 \times 12) = R66\ 443,40$ $\text{Balance} = 235\ 000 \left(1 + \frac{0,11}{12}\right)^{12} - \frac{5\ 536,95 \left[\left(1 + \frac{0,11}{12}\right)^{12} - 1 \right]}{\frac{0,11}{12}}$ $= 192\ 296,17$ $\text{Interest} = (5\ 536,95 \times 12) - (235\ 000 - 192\ 296,17)$ $= 66\ 443,40 - 42\ 703,83$ $= 23\ 739,57$ <p>OR/OF</p>	✓ R66 443,40 ✓ $235\ 000 \left(1 + \frac{0,11}{12}\right)^{12}$ ✓ $\frac{5\ 536,95 \left[\left(1 + \frac{0,11}{12}\right)^{12} - 1 \right]}{\frac{0,11}{12}}$ ✓ R192 296,17 ✓ R42 703,83 ✓ R23 739,57 <p>OR/OF</p>

<p>Total amount paid in first year = $R\ 5\ 536,95 \times 12$ $= R\ 66\ 443,40$</p> <p>Balance on loan after 1 year = P of remaining installments</p> $P = \frac{x[1 - (1 + i)^{-n}]}{i}$ $= \frac{5\ 536,95 \left[1 - \left(1 + \frac{0,11}{12}\right)^{-42}\right]}{\frac{0,11}{12}}$ $= R\ 192\ 296,20$ <p>Amount paid off in the first year: $R\ 235\ 000 - R\ 192\ 296,20 = R\ 42\ 703,80$</p> <p>Amount of interest = $R\ 66\ 443,40 - R\ 42\ 703,80$ $= R\ 23\ 739,60$</p> <p>OR/OF</p> $P = \frac{5536,95 \left[1 - \left(1 + \frac{0,11}{12}\right)^{-12}\right]}{\frac{0,11}{12}}$ $= R\ 62\ 648,18$ <p>$235\ 000 - 62\ 648,18 = R\ 172\ 351,82$</p> <p>After 12 months, money owed on house is</p> $172\ 351,82 \left(1 + \frac{0,11}{12}\right)^{12}$ $= 192\ 296,17$ <p>Amount paid after 12 months is $5\ 536,95 \times 12 = R\ 66\ 443,40$</p> <p>Amount of interest paid: $R\ 66\ 443,40 - (235\ 000 - 192\ 296,17)$ $= R\ 23\ 739,57$</p>	<p>✓ R66 443,40</p> <p>✓ $n = -42$</p> <p>✓ substitution into correct formula</p> <p>✓ R192 296,20</p> <p>✓ R42 703,80</p> <p>✓ R23 739,60</p> <p>OR/OF</p> <p>✓ R62 648,18</p> <p>✓ R172 351,82</p> <p>✓ R192 296,17</p> <p>✓ R66 443,40</p> <p>✓ $235\ 000 - 192\ 296,17$</p> <p>✓ R23 739,57</p> <p style="text-align: right;">(6) [15]</p>
--	---

QUESTION/VRAAG 7

<p>7.1</p> $ \begin{aligned} f(x+h) &= 2(x+h)^2 - (x+h) \\ &= 2(x^2 + 2xh + h^2) - x - h \\ &= 2x^2 + 4xh + 2h^2 - x - h \end{aligned} $ $ \begin{aligned} f(x+h) - f(x) &= 2x^2 + 4xh + 2h^2 - x - h - 2x^2 + x \\ &= 4xh + 2h^2 - h \end{aligned} $ $ \begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{4xh + 2h^2 - h}{h} \\ &= \lim_{h \rightarrow 0} \frac{h(4x + 2h - 1)}{h} \\ &= \lim_{h \rightarrow 0} (4x + 2h - 1) \\ &= 4x - 1 \end{aligned} $ <p>OR/OF</p> $ \begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{2(x+h)^2 - (x+h) - (2x^2 - x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{2x^2 + 4xh + 2h^2 - x - h - 2x^2 + x}{h} \\ &= \lim_{h \rightarrow 0} \frac{4xh + 2h^2 - h}{h} \\ &= \lim_{h \rightarrow 0} \frac{h(4x + 2h - 1)}{h} \\ &= \lim_{h \rightarrow 0} (4x + 2h - 1) \\ &= 4x - 1 \end{aligned} $	<p>✓ $2x^2 + 4xh + 2h^2 - x - h$</p> <p>✓ $4xh + 2h^2 - h$</p> <p>✓ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$</p> <p>✓ subst. into formula</p> <p>✓ $\lim_{h \rightarrow 0} (4x + 2h - 1)$</p> <p>✓ $4x - 1$</p> <p>OR/OF</p> <p>✓ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$</p> <p>✓ subst. into formula</p> <p>✓ $2x^2 + 4xh + 2h^2 - x - h$</p> <p>✓ $4xh + 2h^2 - h$</p> <p>✓ $\lim_{h \rightarrow 0} (4x + 2h - 1)$</p> <p>✓ $4x - 1$</p> <p>(6)</p>
<p>7.2.1</p> $ \begin{aligned} D_x[(x+1)(3x-7)] \\ &= D_x(3x^2 - 4x - 7) \\ &= 6x - 4 \end{aligned} $	<p>✓ $3x^2 - 4x - 7$</p> <p>✓ $6x - 4$</p> <p>(2)</p>
<p>7.2.2</p> $ \begin{aligned} y &= \sqrt{x^3} - \frac{5}{x} + \frac{1}{2}\pi \\ y &= x^{\frac{3}{2}} - 5x^{-1} + \frac{1}{2}\pi \\ \frac{dy}{dx} &= \frac{3}{2}x^{\frac{1}{2}} + 5x^{-2} \end{aligned} $	<p>✓ $x^{\frac{3}{2}} - 5x^{-1}$</p> <p>✓ $\frac{3}{2}x^{\frac{1}{2}}$</p> <p>✓ $+ 5x^{-2}$</p> <p>✓ derivative of $\frac{1}{2}\pi$ is 0</p> <p>(4)</p> <p>[12]</p>

QUESTION/VRAAG 8

8.1	$f(x) = x^3 - 6x^2 + 9x$ $f'(x) = 3x^2 - 12x + 9$ $f''(x) = 6x - 12 = 0$ $x = 2$ $f''(0) = 6(0) - 12 = -12$ $f''(3) = 6(3) - 12 = 6$ <p style="text-align: center;">$\leftarrow f''(x) < \underset{2}{ } f''(x) > 0 \rightarrow$</p> <p>Point of inflection at $x = 2$</p>	✓ $x^3 - 6x^2 + 9x$ ✓ $3x^2 - 12x + 9$ ✓ $6x - 12$ ✓ $6x - 12 = 0$ ✓ explanation (5)
8.2		✓ shape ✓ (0 ; 0) ✓ (3 ; 0) as TP ✓ (1 ; 4) (4)
8.3	f concave up for $x > 2$ $y = -f(x)$ will be concave down for $x > 2$	✓ ✓ $x > 2$ (2)
8.4.1	(3; 7)	✓ 3 ✓ 7 (2)
8.4.2	Do not agree with Claire as her statement is incorrect. Between $x = 1$ and $x = 3$ the graph of f is decreasing. Therefore at $x = 2$ the gradient will have a negative value. <i>Stem nie saam met Claire nie, want haar stelling is verkeerd.</i> <i>Die grafiek van f is dalend/afnemend tussen $x = 1$ en $x = 3$.</i> <i>By $x = 2$ moet die gradiënt dus 'n negatiewe waarde hê.</i> <p style="text-align: center;">OR/OF</p> $f'(2) = 3(2)^2 - 12(2) + 9$ $= -3$ $\neq 1$	✓ no ✓ justification (2) [15]

QUESTION/VRAAG 9

$y = x^2 + 2$ $P(x; x^2 + 2)$ $B(0; 3)$ $\begin{aligned} PB^2 &= (x - 0)^2 + (x^2 + 2 - 3)^2 \\ &= x^2 + x^4 - 2x^2 + 1 \\ &= x^4 - x^2 + 1 \end{aligned}$ <p>PB will be a minimum if PB^2 is a minimum</p> $\begin{aligned} \frac{d(PB^2)}{dx} &= 4x^3 - 2x \\ 4x^3 - 2x &= 0 \\ x(2x^2 - 1) &= 0 \\ x = 0 \text{ or } x^2 &= \frac{1}{2} \\ x &= \frac{1}{\sqrt{2}} \end{aligned}$ $\begin{aligned} PB^2 &= \left(\frac{1}{\sqrt{2}}\right)^4 - \left(\frac{1}{\sqrt{2}}\right)^2 + 1 \\ &= \frac{1}{4} - \frac{1}{2} + 1 \\ &= \frac{3}{4} \\ PB &= \frac{\sqrt{3}}{2} = 0,87 \end{aligned}$ <p>OR/OF</p>	$\checkmark (x - 0)^2 + (x^2 + 2 - 3)^2$ $\checkmark x^4 - x^2 + 1$ $\checkmark 4x^3 - 2x$ $\checkmark \frac{d(PB^2)}{dx} = 0$ $\checkmark x = \frac{1}{\sqrt{2}}$ $\checkmark PB^2 = \left(\frac{1}{\sqrt{2}}\right)^4 - \left(\frac{1}{\sqrt{2}}\right)^2 + 1$ \checkmark answer <p>OR/OF</p>
---	---

	<p>Gradient of tangent to curve = $2x$</p> <p>Gradient of line joining B and the curve = $\frac{x^2 + 2 - 3}{x - 0}$</p> $= \frac{x^2 - 1}{x}$ <p>Shortest distance will be where tangent to curve is perpendicular to the line joining P and the curve.</p> $\frac{x^2 - 1}{x} = -\frac{1}{2x}$ $2x(x^2 - 1) = -x$ $2x^3 - 2x = 0$ $x(2x^2 - 1) = 0$ $x = 0 \quad \text{or} \quad x^2 = \frac{1}{2}$ $x = \frac{1}{\sqrt{2}}$ $\text{PB}^2 = \left(\frac{1}{\sqrt{2}}\right)^4 - \left(\frac{1}{\sqrt{2}}\right)^2 + 1$ $= \frac{1}{4} - \frac{1}{2} + 1$ $= \frac{3}{4}$ $\text{PB} = \frac{\sqrt{3}}{2} = 0,87$	<p>$\checkmark = 2x$</p> <p>$\checkmark = \frac{x^2 - 1}{x}$</p> <p>$\checkmark \frac{x^2 - 1}{x} = -\frac{1}{2x}$</p> <p>$\checkmark 2x^3 - 2x = 0$</p> <p>$\checkmark x = \frac{1}{\sqrt{2}}$</p> <p>$\checkmark \text{PB}^2 = \left(\frac{1}{\sqrt{2}}\right)^4 - \left(\frac{1}{\sqrt{2}}\right)^2 + 1$</p> <p>$\checkmark \text{answer}$</p>
	<p>OR/OF</p> <p>$P(k; k^2 + 2)$ and $B(0; 3)$</p> <p>$BP \perp$ tangent passing through $y = x^2 + 2$ at P.</p> $m_{\text{tangent at } P} = 2k$ $m_{BP} = -\frac{1}{2k}$ <p>Equation of BP: $y = \left(-\frac{1}{2k}\right)x + 3$</p> $y_P = \left(-\frac{1}{2k}\right)(k) + 3 = 2,5$ $\Rightarrow k^2 + 2 = 2,5 \text{ and so } k = \sqrt{0,5} \text{ and } P(\sqrt{0,5}; 2,5)$ $\text{BP} = \sqrt{(\sqrt{0,5} - 0)^2 + (2,5 - 3)^2} = \sqrt{\frac{3}{4}} = \frac{\sqrt{3}}{2} = 0,87$	<p>OR/OF</p> <p>$\checkmark P(k; k^2 + 2)$</p> <p>$\checkmark m_{\text{tangent at } P} = 2k$</p> <p>$\checkmark m_{BP} = -\frac{1}{2k}$</p> <p>$\checkmark y = \left(-\frac{1}{2k}\right)x + 3$</p> <p>$\checkmark \text{value of } y \text{ at P}$</p> <p>$\checkmark \text{value of } k$</p> <p>$\checkmark \text{answer}$</p>

QUESTION/VRAAG 10

10.1	<p style="text-align: center;">$n(S) = 100$</p>	<p>8 values need to be placed in correct position:</p> <p>2 or 3 correct: 1 mark 4 or 5 correct: 2 marks 6 or 7 correct: 3 marks 8 correct: 4 marks</p>
10.2	$(49 - x) + x + 8 + 4 + 5 + 2 + (60 - x) + 14 = 100$ $-x + 142 = 100$ $x = 42$	<p>✓ setting up equation</p> <p>✓ answer</p>
10.3	$P(\text{use only one application}) = \frac{7 + 2 + 18}{100}$ $= \frac{27}{100} \text{ or } 27\%$	<p>✓ $\frac{7 + 2 + 18}{100}$</p> <p>✓ answer</p>

QUESTION/VRAAG 11

11.1	$5 \times 5 \times 10 \times 9$ $= 2250$	<p>✓ 5×5 ✓ 10×9 ✓ 2250</p>																								
11.2	<table border="1" style="margin-bottom: 10px; width: 100%;"> <thead> <tr> <th>No of digits used</th> <th>Letters</th> <th>Digits</th> <th>Total</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>5×5</td> <td>10</td> <td>250</td> </tr> <tr> <td>2</td> <td>5×5</td> <td>10×9</td> <td>2 250</td> </tr> <tr> <td>3</td> <td>5×5</td> <td>$10 \times 9 \times 8$</td> <td>18 000</td> </tr> <tr> <td>4</td> <td>5×5</td> <td>$10 \times 9 \times 8 \times 7$</td> <td>126 000</td> </tr> <tr> <td>5</td> <td>5×5</td> <td>$10 \times 9 \times 8 \times 7 \times 6$</td> <td>756 000</td> </tr> </tbody> </table> <p>Codes of two letters and five digits will ensure unique numbers for 700 000 clients.</p>	No of digits used	Letters	Digits	Total	1	5×5	10	250	2	5×5	10×9	2 250	3	5×5	$10 \times 9 \times 8$	18 000	4	5×5	$10 \times 9 \times 8 \times 7$	126 000	5	5×5	$10 \times 9 \times 8 \times 7 \times 6$	756 000	<p>✓ $5 \times 5 \times 10 \times 9 \times 8 \times 7 \times 6$ ✓✓ five digits</p>
No of digits used	Letters	Digits	Total																							
1	5×5	10	250																							
2	5×5	10×9	2 250																							
3	5×5	$10 \times 9 \times 8$	18 000																							
4	5×5	$10 \times 9 \times 8 \times 7$	126 000																							
5	5×5	$10 \times 9 \times 8 \times 7 \times 6$	756 000																							

TOTAL/TOTAAL: 150